

D37SS64RIB





RM 97 | GDU 2360 SS-RIB AVALIABLE RIB: YES

Management & Positioning

- Consistent high yield potential for maturity
- Strong emergence and excellent early vigor
- Semi-flex girthy ear with high test weight grain
- Good tolerance for eyespot, common rust, and northern leaf blight, average for Goss's wilt
- Medium-tall stature

Agronomic Ratings

EMERGENCE
SEEDLING VIGOR
STALK RATING
ROOT RATING
GREENSNAP SCORE
DROUGHT
STAYGREEN
TEST WEIGHT
DRYDOWN
GOSS'S WILT

Agronomic Traits					
Plant Height	Medium-Tall	Kernel Rows	16-18		
Ear Height	Medium	Cob Color	Red		
Flowering		Kernel Texture	Med-Hard		
Leaf Habit	Semi-Upright	Kernel Depth	Medium		
Ear Flex	Semi-Flex	Husk Coverage	Short		
Ear Type	Med-Girth	Shank Length	Medium		

Trait Versions Available

CONV - NONE | D37VC64RIB

Precis	sion Placeme	nt™ Managemeı	nt
Planting Date		Soils	
Early	HR	Clay Loams	R
Late	R	Sandy	R
Variable Planting Populations		Silt Loam	HR
With Yield Zone		Peat	R
Low	24-30,000	Compacted	N
Moderate	28-34,000	Poorly Drained	N
High	34-40,000	Drought Prone	N
Very High	38-42,000	High pH	R
Dryland <20	N	Fertility	
Population=(Yield Goal/7.5)*1000		Nitrogen	
Water Managemer	nt	Low	N
Full Irrigation	HR	Med	R
Limited	HR	High	HR
Dryland	HR	Post Application	
Crop Rotation		Herbicide	Caution: SU
Corn/Soybeans	HR	Fungicide	Positive
Continue Corn	w/Fungicide	LPI Nutritional	Excellent
Tillage		Herbicide Resistance	Glyph / Gluf
Conventional	HR	Harvest Schedule	
Minimum	HR	Early	HR
Ridge-Till	HR	Late	R
No-Till	HR		
Soil Productivity		Forage / Silage Quality	
Low	R	Silage Select	N
Moderate	HR	Dual Purpose	R
High	HR		

Disease Tolerance Ratings					
n/a					
6					
8					
7					
r					

Plant with These Hybrids for Diversity

D35SS58 | D37SS60

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Testing not complete. Herbicide abbreviations: GR=Growth Regulator, PI=Pigment Inhibitor, SU=Sulfonylurea. Yield zones based upon yield goals in field.

Actual ratings based on best current information available and may be affected by changing environmental and management conditions.