

S009XF33





XTFlex 0.09 RM

No-Till

Management & Positioning

- Late group 00 XtendFlex® release is ideal for productive environments in its maturity zone
- Very good tolerance for iron deficiency chlorosis and resistance for brown stem rot
- Medium height plant with good lateral branching and excellent standability
- Rps1k gene for Phytophthora root rot and above average Sclerotinia white mold tolerance

Agronomic Ratings

| EMERGENCE |
|---------------------------|
| STANDABILITY |
| STRESS TOLERANCE |
| SHATTER RESISTANCE |
| PHYTOPHTHORA FIELD TOL. |
| SUDDEN DEATH SYNDROME |
| IRON DEFICIENCY CHLOROSIS |
| SCLEROTINIA WHITE MOLD |
| BROWN STEM ROT |
| |

Phytophthora Field Tolerance

- Score designates reaction to Phytophthora sojae Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to Phytophthora sojae Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

Phytophthora Gene Resistance

| s= | Susceptible | or no | specific gene | resistance |
|----|-------------|-------|---------------|------------|
|----|-------------|-------|---------------|------------|

Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44

Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44

Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45

HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

| Precision Placement™ Management | ľ |
|---------------------------------|---|
|---------------------------------|---|

| Row Width | | Soils | |
|-----------------------------|----|--|---------|
| Wide | N | Clay & Clay Loams | R |
| 15-20" | HR | Sands & Sandy Loams Loams & Silt Loam | R HR |
| Drilled | HR | Poorly Drained IDC | N HR |
| Planting Populations | | High pH | R |
| Greater than 190K | HR | | |
| 160-180K | HR | | |
| 130-150K | Ν | | |
| 100/120K | N | | |
| Tillage | | Yield Environment | |
| Conventional | HR | High | HR |
| Minimum | HR | Stable | HR |

| Ag | | | | Trai | 400 |
|----------|---------------------|----------------|-------------|------------------------|-----------------|
| AG | OH | OIII | | | LS |
| Tariffel | Name and Address of | and the latest | CHICAGO III | NAME OF TAXABLE PARTY. | Suctored |

Stress

Double Crop/Delayed Following Soybeans

HR

| Plant Height | Medium | Hilium Color | Black |
|--------------|----------|----------------------|-----------|
| Canopy Type | Mod-Bush | Oil Content | 18.0-19.0 |
| Flower Color | Purple | Protein Content | 32.0-33.0 |
| Pubescence | Tawny | Metribuzin Rating | 7.0 |
| Pod Color | Brown | Chloride Sensitivity | Includer |
| | | | |

Disease Tolerance Ratings

| | | <u>™</u> | |
|--------------------|-----------|------------------------|-------|
| Cyst Nematode | S | PRR Resistance | Rps1k |
| SCN Resistance | n/a | PRR Field Tolerance | 6 |
| Sclerotinia W. Mol | d 7 | Frogeye Leaf Spot | n/a |
| Brown Stem Rot | 9 | Stem Canker | n/a |
| Sudden Death | n/a | Charcoal Rot | n/a |
| IDC | 7 | S Root Knot Nematode | n/a |
| IDC Recovery | Above Avg | Cercospora Leaf Blight | n/a |

Plant with These Varieties

S006XF83 | S01XF43

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data. Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

Actual ratings based on best current information available and may be affected by changing environmental and management conditions.