

S21XF6





XTFlex 2.1 RM

No-Till

Management & Positioning

- Early group II XtendFlex® release features a sound agronomic package
- Rps1c gene for Phytophthora root rot and very good Sclerotinia white mold tolerance
- Good iron deficiency chlorosis tolerance and resistance for brown stem rot
- Medium-tall plant height with moderate canopy type and excellent standability
- Favors placement on better productive loam soil types within its maturity zone of adaptation

Agronomic Ratings

EMERGENCE
STANDABILITY
STRESS TOLERANCE
SHATTER RESISTANCE
PHYTOPHTHORA FIELD TOL.
SUDDEN DEATH SYNDROME
IRON DEFICIENCY CHLOROSIS
SCLEROTINIA WHITE MOLD
BROWN STEM ROT
0 1 2 2 4 5 6 7 0

Phytophthora Field Tolerance

- Score designates reaction to Phytophthora sojae Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to Phytophthora sojae Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

Phytophthora Gene Resistance

S = Susceptible or no specific gene resistance

Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44

Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44

Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45

HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

Precision Placement ™ Management

Row Width		Soils	
Wide	N	Clay & Clay Loams	R
15-20"	HR	Sands & Sandy Loams Loams & Silt Loam	R HR
Drilled	HR	Poorly Drained	N
Planting Populations		IDC	R
		High pH	N
Greater than 190K	R		
160-180K	HR		
130-150K	R		
100/120K	N		
Tillage		Yield Environment	
Conventional	HR	High	HR
Minimum	HR	Stable	HR
· · · · · · · · · · · · · · · · · · ·		Stress	R

10000				-	
Aa	ron	on	nic	Trai	ts
Tables.	of the Party of th	ANNEXOS	NAMES OF TAXABLE PARTY.	INCOME.	Subset

Double Crop/Delayed Following Soybeans

HR

Plant Height	Med-Tall	Hilium Color	Gray
Canopy Type	Moderate	Oil Content	17.0-18.0
Flower Color	Purple	Protein Content	35.0-36.0
Pubescence	Gray	Metribuzin Rating	5.0
Pod Color	Tan	Chloride Sensitivity	n/a

Disease Tolerance Ratings

Cyst Nematode	R3,MR14	PRR Resistance	Rps1c
SCN Resistance	PI88788	PRR Field Tolerance	7
Sclerotinia W. Mold	7	Frogeye Leaf Spot	n/a
Brown Stem Rot	9	Stem Canker	9
Sudden Death	7	Charcoal Rot	6
IDC	7	S Root Knot Nematode	n/a
IDC Recovery	Average	Cercospora Leaf Blight	n/a

Plant with These Varieties

S17XF02 | S19XF62 | S25XF64

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data. Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

Actual ratings based on best current information available and may be affected by changing environmental and management conditions.