




1.4 RM

ENLIST

Management & Positioning

- Enlist E3® soybean introduction features the Peking source of resistance for cyst nematode
- Resistance for brown stem rot and very good field tolerance for Phytophthora root rot
- Above average tolerance scores for iron deficiency chlorosis and Sclerotinia white mold
- Medium plant height with good lateral branching and very good standability
- Favors placement within its maturity zone of adaptation

Agronomic Ratings



Phytophthora Field Tolerance

- Score designates reaction to *Phytophthora sojae* Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to *Phytophthora sojae* Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

Phytophthora Gene Resistance

S = Susceptible or no specific gene resistance
 Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36
 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44
 Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44
 Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45
 HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data.

Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

****Actual ratings based on best current information available and may be affected by changing environmental and management conditions.****

Precision Placement™ Management

Row Width		Soils	
Wide - 36-40"	N	Clay & Clay Loams	HR
Twin or 30"	R	Sands & Sandy Loams	R
15-20"	HR	Loams & Silt Loam	HR
Drilled	HR	Poorly Drained	R
Planting Populations		IDC	R
		High pH	R
Greater than 190K	R		
160-180K	HR		
130-150K	HR		
100/120K	N		

Tillage		Yield Environment	
Conventional	HR	High	HR
Minimum	HR	Stable	HR
No-Till	HR	Stress	HR
		Double Crop/Delayed	R
		Following Soybeans	HR

Agronomic Traits

Plant Height	M	Hilium Color	BL
Canopy Type	MB	Oil Content	19.0-20.0
Flower Color	P	Protein Content	35.0-36.0
Pubescence	LT	Metribuzin Rating	7.0
Pod Color	BR	Chloride Sensitivity	INC

Disease Tolerance Ratings

Cyst Nematode	R1,R3,MR5	PRR Resistance	Rps1k
SCN Resistance	Peking	PRR Field Tolerance	7
Sclerotinia W. Mold	7	Frogeye Leaf Spot	n/a
Brown Stem Rot	8	Stem Canker	9
Sudden Death	7	Charcoal Rot	6
IDC	6	S Root Knot Nematode	2
IDC Recovery	Average	Cercospora Leaf Blight	7.0

Plant with These Varieties

S09EN26 | S12EN72 | S16EN42