





HR

HR

R

ENLIST 2.0 RM

Minimum

No-Till

Management & Positioning

- Enlist E3® soybean features the Peking source of resistance for soybean cyst nematode
- Excellent tolerance for Sclerotinia white mold with good iron deficiency chlorosis tolerance
- Medium tall plant height, moderate canopy type and very strong standability scores
- Rps3a gene for Phytophthora root rot and resistance for brown stem rot

Agronomic Ratings

EMERGENCE
STANDABILITY
STRESS TOLERANCE
SHATTER RESISTANCE
PHYTOPHTHORA FIELD TOL.
SUDDEN DEATH SYNDROME
IRON DEFICIENCY CHLOROSIS
SCLEROTINIA WHITE MOLD
BROWN STEM ROT

Phytophthora Field Tolerance

- Score designates reaction to Phytophthora sojae Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to Phytophthora sojae Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

Phytophthora Gene Resistance

S	=	Susceptible	or	no	specific	gene	resistance
---	---	-------------	----	----	----------	------	------------

Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44

Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44

Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45

HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

Precision Placement™ Management

		1873	
Row Width		Soils	
Wide	N	Clay & Clay Loams	HR
15-20"	HR	Sands & Sandy Loams Loams & Silt Loam	R HR
Drilled	HR	Poorly Drained	N
Planting Populations		IDC High pH	HR R
Greater than 190K	R		
160-180K	HR		
130-150K	HR		
100/120K	N		
Tillage		Yield Environment	
Conventional	HR	High	HR
Minimum	ЦΒ	Stable	HR

Stress

Double Crop/Delayed

Following Soybeans

HR

HR

Agronomic Traits					
Plant Height	M/T	Hilium Color	IB		
Canopy Type	M	Oil Content	n/a		
Flower Color	Р	Protein Content	n/a		
Pubescence	G	Metribuzin Rating	n/a		
Pod Color	TN	Chloride Sensitivity	INC		

Disease Tolerance Ratings

•					
Cyst Nematode	MR1,R3	PRR Resistance	Rps3a		
SCN Resistance	Peking	PRR Field Tolerance	7		
Sclerotinia W. Mold	7	Frogeye Leaf Spot	n/a		
Brown Stem Rot	9	Stem Canker	9		
Sudden Death	6	Charcoal Rot	7		
IDC	7	S Root Knot Nematode	2		
IDC Recovery	Average	Cercospora Leaf Blight	n/a		
		1			

Plant with These Varieties

S18EN35 | S20EN84 | S21EN81 | S23EN05

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data. Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

Actual ratings based on best current information available and may be affected by changing environmental and management conditions.

2025 Loveland Products, Inc. All Rights Reserved. Dyna-Gro is a registered trademark of Loveland Products, Inc. All other trademarks are the property of their respective owners.

The transgenic soybean event in Enlist E3® soybeans is jointly developed and owned by Corteva Agriscience LLC & M.S. Technologies, LLC. Enlist products contain the Enlist trait provides crop safety for use of labeled over-the-top applications of glyphosate, glufosinate & 2,4-D herbicides featuring Colex-D® technology when applied according to label directions. 2,4-D products that do not contain Colex-D® technology when applied according to label directions. technology are not authorized for use with Enlist products. Enlist, Enlist E3, the Enlist E3 logo and Colex-D are trademarks of Corteva Agriscience and its affiliated companies. For complete soybean stewardship and trait legal statements, please refer to the Dyna-Gro® Product Guide.