





3.5 RM **ENLIST/STS** 

## **Management & Positioning**

- Mid-group III Enlist E3® soybean release with STS® with exceptional performance in productive environments
- Rps1c gene with good field tolerance for Phytophthora root rot and resistance for cyst nematode
- Resistance for stem canker and above average tolerance for sudden death syndrome
- Medium compact plant height with moderate lateral branching and excellent standability scores

# **Agronomic Ratings**

EMERGENCE
STANDABILITY
STRESS TOLERANCE
SHATTER RESISTANCE
PHYTOPHTHORA FIELD TOL.
SUDDEN DEATH SYNDROME
IRON DEFICIENCY CHLOROSIS
SCLEROTINIA WHITE MOLD
BROWN STEM ROT

# Phytophthora Field Tolerance

- Score designates reaction to Phytophthora sojae Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to Phytophthora sojae Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

#### Phytophthora Gene Resistance

s=	Susceptible	or no speci	fic gene	resistance
----	-------------	-------------	----------	------------

Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44

Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44

Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45

HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

# Precision Placement™ Management

Row Width		Soils	
Wide	N	Clay & Clay Loams	HR
Twin or 30	HR	Sands & Sandy Loams	R
15-20"	HR	Loams & Silt Loam	HR
Drilled	HR	Poorly Drained	N
Planting Populations		IDC High pH	R N
Greater than 190K	N	1g p	
160-180K	HR		
130-150K	HR		
100/120K	R		
Tillage		Yield Environment	
Conventional	HR	High	HR
Minimum	ЦΒ	Stable	HR

Minimum	HR	0.00.0	
<del></del>		Stress	HR
No-Till	HR	Double Crop/Delayed	R
		Double Crop/Delayed Following Soybeans	R

Agronomic Traits			
Plant Height	М	Hilium Color	BL
Canopy Type	M	Oil Content	18.0-19.0
Flower Color	W	Protein Content	33.0-34.0
Pubescence	LT	Metribuzin Rating	7.0
Pod Color	BR	Chloride Sensitivity	INC
	LI BR	Chloride Sensitivity	_

#### **Disease Tolerance Ratings**

•			
Cyst Nematode	R3	PRR Resistance	Rps1c
SCN Resistance	PI88788	PRR Field Tolerance	7
Sclerotinia W. Mold	5	Frogeye Leaf Spot	n/a
Brown Stem Rot	3	Stem Canker	9
Sudden Death	7	Charcoal Rot	6
IDC	6	S Root Knot Nematode	2
IDC Recovery	Average	Cercospora Leaf Blight	n/a

### **Plant with These Varieties**

S33ES76 | S36EN36 | S37ES52 | S38EN75

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data. Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

\*\*Actual ratings based on best current information available and may be affected by changing environmental and management conditions.\*\*