

S68XF16S





HR

HR

6.8 RM XTFlex/STS

No-Till

Management & Positioning

- Determinate late group VI XtendFlex® & STS® introduction that is ideal for lighter soils and stress environments
- Intermediate tolerance for Southern root knot nematode and excluder for chloride salts
- Taller plant height with nice lateral branching for wider row widths and average standability scores
- A planned fungicide program may be warranted for frogeye leaf spot control

Agronomic Ratings

EMERGENCE
STANDABILITY
STRESS TOLERANCE
SHATTER RESISTANCE
PHYTOPHTHORA FIELD TOL.
SUDDEN DEATH SYNDROME
STEM CANKER
FROGEYE LEAF SPOT
S. ROOT KNOT NEMATODE

Phytophthora Field Tolerance

- Score designates reaction to Phytophthora sojae Race 25 for commercial genes Rps1a, Rps1c and Rps1k.
- Score designates reaction to Phytophthora sojae Race 30 for commercial gene Rps3a. Score also based upon in-field observations.
- Phytophthora Field Tolerance scores are important for races of Phytophthora not covered by specific genes of resistance.

Phytophthora Gene Resistance

s=	Susceptible	or no	specific	gene	resistance
----	-------------	-------	----------	------	------------

Rps1a = Denotes resistance to Races 1, 2, 10, 11, 13-18, 24, 26, 27, 31, 32 and 36 Rps1c = Denotes resistance to Races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, 32, 34, 36, 41, 42 and 44

Rps1k = Denotes resistance to Races 1-11, 13-15, 17, 18, 21-24, 26, 36, 37 and 42-44

Rps3a = Denotes resistance to Races 1-5, 8, 9, 11, 13, 14, 16, 18, 23, 25, 28, 29, 31-35, 40 and 43-45

HRps = Denotes Heterozygous resistance (partial resistance) to the specific gene noted

Precision Placement™ Management

Row Width		Soils	
Wide	HR	Clay & Clay Loams	HR
Twin or 30	HR	Sands & Sandy Loams	HR
15-20"	HR	Loams & Silt Loam	R
Drilled	R	Poorly Drained	N
Planting Populations		IDC High pH	N N
Greater than 190K	Ν		
160-180K	Ν		
130-150K	HR		
100/120K	HR		
Tillage		Yield Environment	
Conventional	HR	High	HR
Minimum	HR	Stable	HR

Stress

Double Crop/Delayed

Following Soybeans

HR

Agronomic Traits						
Plant Height	Т	Hilium Color	BL			
Canopy Type	В	Oil Content	18.0-19.0			
Flower Color	Р	Protein Content	35.0-36.0			
Pubescence	Т	Metribuzin Rating	5.0			
Pod Color	BR	Chloride Sensitivity	EXC			

Disease Tolerance Ratings

Cyst Nematode	S	PRR Resistance	HRps1c	
SCN Resistance	n/a	PRR Field Tolerance	6	
Sclerotinia W. Mold	n/a	Frogeye Leaf Spot	5	
Brown Stem Rot	n/a	Stem Canker	7	
Sudden Death	n/a	Charcoal Rot	n/a	
IDC	n/a	S Root Knot Nematode	5	
IDC Recovery	n/a	Cercospora Leaf Blight	n/a	

Plant with These Varieties

S63XF56 | S67XF34S | S71XF93

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Insufficient Data. Soybean Cyst Nematode: R=Resistant, MR=Moderately Resistant, S=Susceptible, # Denotes race number for resistance.

Actual ratings based on best current information available and may be affected by changing environmental and management conditions.